



Volume 6, Issue 1, 2023

Peer-reviewed, open-access journal dedicated to publishing high-quality research on sustainable technology and infrastructure in developing countries.

<https://research.tensorgate.org/index.php/tjstidc>

ACCESS TO MENTAL HEALTHCARE SERVICES IN DEVELOPING COUNTRIES: A REVIEW OF CHALLENGES AND TECH-INTERVENTIONS

Mohammed Irfan Jawad

Mehran University of Engineering and Technology, Jamshoro, Pakistan

ABSTRACT

This study explores the challenges and interventions related to accessing mental healthcare services in developing countries. The findings reveal that stigma, lack of resources, low awareness, and cultural barriers are significant challenges that prevent individuals from accessing the necessary care. Stigma associated with mental illness remains a significant barrier to accessing care in many developing countries. Lack of resources and infrastructure, including a shortage of mental health professionals, medication, and treatment facilities, also pose a challenge. Additionally, a lack of awareness and understanding about mental health, as well as cultural beliefs and practices, can delay seeking help and limit access to appropriate care. Community-based interventions, task-sharing, telemedicine, and advocacy are identified as effective interventions to address these challenges. Community-based interventions can increase access to care by providing education programs, outreach services, and community-based clinics. Task-sharing, which involves training non-specialist healthcare providers to offer basic mental healthcare services, can be an effective way to provide care in areas where mental health professionals are scarce. Telemedicine can also connect patients with mental health professionals remotely, particularly in remote or underserved areas. Advocacy efforts, including public awareness campaigns, media outreach, and lobbying for policy changes, can reduce stigma and increase awareness about mental illness. This study highlights the need for increased access to mental healthcare services in developing countries. Interventions such as community-based interventions, task-sharing, telemedicine, and advocacy can help overcome the challenges of stigma, lack of resources, low awareness, and cultural barriers. These interventions can improve access to care and ultimately improve the lives of individuals living with mental illness in developing countries.

Keywords: *Mental healthcare services, Developing countries, Stigma, Task-sharing, Telemedicine*

INTRODUCTION

Mental healthcare refers to the provision of services and support to individuals who are experiencing mental health challenges. This includes a range of services, from diagnosis and treatment to ongoing support and recovery. Mental healthcare is an important aspect of overall healthcare, as mental health issues can have a significant impact on an individual's physical health, social functioning, and overall quality of life.

There are many different types of mental health issues that individuals may experience, ranging from anxiety and depression to more severe conditions such as bipolar disorder, schizophrenia, and personality disorders. Mental health issues can be caused by a variety of factors, including genetics, environmental factors, and life experiences. It is important for mental healthcare providers to be able to diagnose and treat these conditions in a way that is effective and compassionate.



Tensorgate Journal of Sustainable Technology and Infrastructure for Developing Countries

<https://research.tensorgate.org/index.php/tjstidc>

One of the key components of mental healthcare is diagnosis. This involves the assessment of an individual's mental health status, including an evaluation of their symptoms and an assessment of any potential risk factors. Diagnosis is important because it allows mental healthcare providers to determine the most appropriate course of treatment for an individual's particular condition.

Treatment for mental health issues can take many different forms, depending on the nature and severity of the condition. Common treatments include medication, therapy, and lifestyle changes. Medication is often used to manage symptoms of mental health conditions, such as anxiety, depression, and bipolar disorder. Therapy can also be an important part of treatment, as it allows individuals to talk through their experiences and develop coping strategies for managing their symptoms. Lifestyle changes such as exercise, healthy eating, and stress reduction techniques can also be effective in managing mental health issues.

In addition to diagnosis and treatment, mental healthcare also involves ongoing support and recovery. This can include support groups, therapy sessions, and other forms of support that help individuals to maintain their mental health and prevent relapse. Recovery from mental health issues is a process that can take time, and it is important for mental healthcare providers to work with individuals to develop a personalized plan that addresses their specific needs.

Access to mental healthcare is a critical issue that affects millions of individuals worldwide. While mental health issues are common, many individuals do not receive the care they need due to barriers such as stigma, cost, and limited availability of mental healthcare resources. Improving access to mental healthcare is essential to promoting overall health and well-being and reducing the burden of mental illness on individuals, families, and communities.

One of the primary barriers to mental healthcare access is stigma. Stigma is a negative attitude or belief about mental health conditions that can lead to discrimination and prejudice. Many individuals who experience mental health challenges may feel ashamed or embarrassed to seek help due to the stigma associated with mental illness. This can lead to delays in seeking treatment or a reluctance to seek help at all. To address this issue, it is important to educate individuals about mental health and to reduce the stigma associated with mental illness. This can be achieved through public education campaigns, advocacy, and promoting positive representations of mental health conditions in media and popular culture.

Another barrier to mental healthcare access is cost. Mental healthcare services can be expensive, and many individuals may not have access to affordable care. This is especially true for those without health insurance or those whose insurance does not cover mental healthcare. To address this issue, it is important to expand access to affordable mental healthcare services. This can be achieved through the implementation of policies that ensure mental healthcare is covered by insurance plans and the creation of programs that provide low-cost or free mental healthcare services to individuals who cannot afford to pay.

Limited availability of mental healthcare resources is another significant barrier to access. In many areas, mental healthcare resources may be limited, particularly in rural or

underserved communities. This can make it difficult for individuals to access care, particularly if they do not have the means to travel to another location to receive treatment. To address this issue, it is important to invest in mental healthcare infrastructure and resources in underserved areas. This can include the creation of telehealth programs that allow individuals to receive mental healthcare services remotely and the establishment of mental healthcare clinics in underserved areas.

Another important factor affecting mental healthcare access is cultural competence. Mental healthcare providers must be able to provide care that is culturally sensitive and responsive to the needs of diverse populations. This includes individuals from different ethnic and racial backgrounds, individuals with disabilities, and individuals from the LGBTQ+ community. Cultural competence is important because it helps to ensure that individuals receive care that is respectful, responsive, and appropriate to their unique needs and circumstances.

In addition to these barriers, there are also systemic issues that affect mental healthcare access. One of these issues is the shortage of mental healthcare providers. There is a significant shortage of mental healthcare providers in many areas, which can make it difficult for individuals to access care. This shortage is particularly acute in rural areas, where there are fewer mental healthcare providers per capita than in urban areas. To address this issue, it is important to invest in mental healthcare provider training programs and to provide incentives for mental healthcare providers to practice in underserved areas.

Another systemic issue that affects mental healthcare access is the fragmentation of mental healthcare services. Mental healthcare services are often delivered through a fragmented system of care, with different providers and organizations responsible for different aspects of care. This can lead to gaps in care and difficulty navigating the healthcare system. To address this issue, it is important to create integrated mental healthcare systems that provide coordinated and comprehensive care across all aspects of mental healthcare, from diagnosis and treatment to ongoing support and recovery.

Improving access to mental healthcare is an essential step towards promoting overall health and well-being and reducing the burden of mental illness on individuals and communities. By addressing the barriers to mental healthcare access, we can ensure that individuals receive the care they need to manage their mental health challenges and improve their quality of life.

CHALLENGES

Mental health has become an increasingly important topic in recent years, as it affects people of all ages, genders, and backgrounds. While there has been significant progress in many developed countries in addressing mental health issues, many developing countries continue to struggle with providing adequate mental health care to their populations. There are several significant barriers to mental health care in developing countries, including stigma, lack of resources, low awareness, and cultural barriers.

Stigma is one of the most significant barriers to mental health care in developing countries. Mental illness is often stigmatized, and people with mental health problems may be seen as weak, dangerous, or unpredictable. This stigma can lead to discrimination, social exclusion, and a lack of support from family and friends. It can also lead to a reluctance to

seek treatment, as people may be afraid of being judged or ostracized by their communities. Stigma can affect people with any type of mental illness, including depression, anxiety, schizophrenia, and bipolar disorder.

Stigma is particularly prevalent in developing countries, where mental illness is often seen as a sign of weakness or moral failing. In some cultures, mental illness is associated with supernatural or spiritual causes, which can lead to further stigma and discrimination. For example, in some African countries, mental illness is often thought to be caused by witchcraft or possession by evil spirits. This can lead to people with mental illness being ostracized from their communities or even attacked.

Table 1. Barriers to access to mental healthcare	
Factors	Description
Stigma	Mental illness is often stigmatized in many developing countries, making it difficult for individuals to seek treatment.
Lack of Resources	Developing countries often lack the necessary resources and infrastructure to provide mental healthcare services.
Low Awareness	There is often a lack of awareness and understanding about mental health in developing countries, leading to a delay in seeking help and a lack of access to appropriate care.
Cultural Barriers	Cultural beliefs and practices can also be a significant barrier to accessing mental healthcare services in some countries.

Lack of resources is another significant barrier to mental health care in developing countries. Many developing countries lack the necessary resources and infrastructure to provide mental healthcare services. This includes a shortage of mental health professionals, medication, and treatment facilities. In many countries, there are only a handful of psychiatrists or psychologists, and they may be concentrated in urban areas, leaving rural communities without access to mental health care.

Even when mental health professionals are available, they may not have the necessary resources to provide adequate care. For example, in some countries, mental health professionals may not have access to the latest medications or treatments, or they may not have the necessary training to use them effectively. Additionally, mental health facilities may be poorly equipped, with outdated or inadequate facilities and equipment.

Low awareness is also a significant barrier to mental health care in developing countries. There is often a lack of awareness and understanding about mental health, which can lead to a delay in seeking help and a lack of access to appropriate care. Many people in developing countries may not recognize the symptoms of mental illness or may not understand that it is a medical condition that requires treatment.

Low awareness is often compounded by a lack of education about mental health. In many developing countries, mental health is not taught in schools, and there may be few public education campaigns or awareness-raising initiatives. This lack of education can lead to misconceptions about mental health and a lack of understanding about the importance of seeking treatment.

Cultural barriers are also a significant barrier to accessing mental healthcare services in developing countries. Cultural beliefs and practices can impact how mental health is perceived and treated in different cultures. Some cultures may not recognize mental illness as a medical condition and may rely on traditional healers or spiritual leaders for treatment. This can make it difficult for people with mental illness to access evidence-based treatment and may lead to delays in seeking help.

Cultural barriers can also affect how mental health is perceived within families and communities. For example, in some cultures, mental illness is seen as a personal failure or a sign of weakness, which can lead to stigma and discrimination. Additionally, some cultures may not have words or concepts for certain mental health conditions, making it difficult for people to express their symptoms or for mental health professionals to provide accurate diagnoses. Mental health care in developing countries is impacted by a range of barriers, including stigma, lack of resources, low awareness, and cultural barriers. These barriers can prevent people from accessing the care they need, leading to a worsening of their mental health conditions and poorer quality of life.

INTERVENTIONS

Community-based interventions, task-sharing, telemedicine, and advocacy are strategies that have been used to increase access to mental healthcare services in developing countries. Community-based interventions refer to programs or services that are developed and delivered within a particular community. These interventions aim to increase access to mental healthcare services by providing education programs, outreach services, and community-based clinics. The primary goal of community-based interventions is to improve mental health outcomes and reduce the stigma associated with mental illness.

Interventions	Description
Community-Based Interventions	Community-based interventions, such as education programs and outreach services, can increase access to mental healthcare services in developing countries.
Task-Sharing	Task-sharing involves training non-specialist healthcare providers to provide basic mental healthcare services, which can be an effective way to increase access to care.
Telemedicine	Telemedicine uses technology to provide mental healthcare services to individuals in remote or underserved areas.
Advocacy	Advocacy efforts can increase awareness and reduce stigma around mental illness through public awareness campaigns, media outreach, and policy changes.

Community-based interventions are effective in developing countries because they address the specific needs and circumstances of the community. For example, community-based clinics can be set up in remote areas where individuals may not have access to mental healthcare services. Outreach services can be used to provide mental healthcare services to individuals who are unable to travel to a clinic. Education programs can be used to increase awareness about mental illness and reduce stigma within the community.

One example of a successful community-based intervention is the Friendship Bench program in Zimbabwe. This program trains lay health workers to provide counseling

services to individuals who are experiencing depression. The program has been successful in increasing access to mental healthcare services and improving mental health outcomes in Zimbabwe. Similar programs have been implemented in other developing countries, including India and Ethiopia.

Task-sharing involves training non-specialist healthcare providers to provide basic mental healthcare services. This approach can be an effective way to increase access to care in areas where there is a shortage of mental health professionals. Non-specialist healthcare providers can include community health workers, nurses, and general practitioners.

Task-sharing is effective in developing countries because it allows for the delivery of mental healthcare services in areas where there is a shortage of mental health professionals. It also increases the capacity of the healthcare system to provide mental healthcare services. However, task-sharing also has limitations, as non-specialist healthcare providers may lack the necessary training and resources to provide high-quality mental healthcare services.

One example of a successful task-sharing program is the Mental Health Gap Action Program (mhGAP) in Uganda. This program trains primary healthcare providers to diagnose and treat common mental health conditions. The program has been successful in increasing access to mental healthcare services and improving mental health outcomes in Uganda. Similar programs have been implemented in other developing countries, including Kenya and Haiti.

Telemedicine involves using technology to provide healthcare services to patients remotely. Telemedicine can be an effective way to provide mental healthcare services to individuals in remote or underserved areas. This approach uses technology to connect patients with mental health professionals remotely, including video conferencing, telephone consultations, and mobile applications.

Telemedicine is effective in developing countries because it can be used to provide mental healthcare services to individuals in remote or underserved areas. It also increases access to mental healthcare services for individuals who may not have the resources or ability to travel to a clinic. However, telemedicine also has limitations, as it requires access to reliable technology and may not be feasible in areas with limited infrastructure.

One example of a successful telemedicine program is the StrongMinds program in Uganda. This program uses a mobile application to provide group therapy sessions to individuals who are experiencing depression. The program has been successful in increasing access to mental healthcare services and improving mental health outcomes in Uganda. Similar programs have been implemented in other developing countries, including India and South Africa.

Advocacy refers to efforts to increase awareness and reduce stigma around mental illness. Advocacy efforts can include public awareness campaigns, media outreach, and lobbying for policy changes to improve access to care. Advocacy is important in developing countries because mental illness is often stigmatized, which can prevent individuals from seeking treatment. Advocacy efforts can help to reduce the stigma associated with mental illness and increase awareness about the importance of mental healthcare services.

One example of a successful advocacy program is the BasicNeeds program in India. This program uses a community-based approach to increase awareness about mental illness and improve access to mental healthcare services. The program has been successful in reducing stigma and increasing access to mental healthcare services in India. Similar programs have been implemented in other developing countries, including Kenya and Ghana. Community-based interventions, task-sharing, telemedicine, and advocacy are strategies that have been used to increase access to mental healthcare services in developing countries. These strategies address the specific needs and circumstances of the community and increase the capacity of the healthcare system to provide mental healthcare services. However, each approach has limitations, and a combination of strategies may be necessary to address the significant burden of untreated mental illness in developing countries. Addressing these issues requires significant investments in mental healthcare infrastructure, training of mental health professionals, and policy changes to prioritize mental healthcare services. However, community-based interventions, task-sharing, telemedicine, and advocacy are important steps in increasing access to mental healthcare services in developing countries and reducing the burden of untreated mental illness.

AI and ML have the potential to significantly improve access to mental healthcare services in developing countries. Firstly, AI and ML can assist in the early detection of mental health disorders. With the help of machine learning algorithms, patterns and trends in data can be analyzed to identify individuals who are at high risk of developing mental health disorders. This can help in early intervention and prevention, which can be critical in reducing the burden of mental health disorders.

Secondly, AI and ML can help in the development of personalized treatment plans. Mental health disorders are complex and vary from person to person. With the help of AI and ML algorithms, individual patient data can be analyzed to develop personalized treatment plans. This can help in providing more effective and efficient treatment, which can be crucial in developing countries where access to mental healthcare services is limited.

Thirdly, AI and ML can assist in improving the quality of care provided to patients. With the help of AI and ML algorithms, healthcare professionals can analyze patient data to identify areas of improvement in the treatment plan. This can help in providing more personalized and effective care, which can ultimately improve patient outcomes.

Challenges of AI and ML in Mental Healthcare Services: Despite the potential benefits, there are also several challenges associated with the use of AI and ML in mental healthcare services in developing countries. Firstly, the lack of data infrastructure and resources is a major challenge. Developing countries often have limited resources to collect and analyze data, which can make it difficult to implement AI and ML solutions.

Secondly, there is a lack of trained professionals in developing countries who can effectively use AI and ML tools. The success of AI and ML in mental healthcare services depends on the availability of trained professionals who can effectively use these tools. However, developing countries often have a shortage of trained professionals, which can hinder the implementation of AI and ML solutions.

Thirdly, there is a lack of awareness and acceptance of AI and ML among patients and healthcare professionals in developing countries. Patients and healthcare professionals may

be skeptical of using AI and ML in mental healthcare services, which can limit the implementation of these solutions.

Implications of AI and ML for Developing Countries: The use of AI and ML in mental healthcare services has significant implications for developing countries. Firstly, the implementation of AI and ML solutions can help in overcoming the shortage of trained professionals in developing countries. AI and ML tools can assist healthcare professionals in providing more effective and efficient care, which can ultimately improve patient outcomes. Secondly, the use of AI and ML can help in reducing the burden of mental health disorders in developing countries. Early detection and prevention of mental health disorders can help in reducing the burden of these disorders, which can have significant economic and social benefits. Thirdly, the use of AI and ML can help in improving the quality of care provided to patients in developing countries. With the help of AI and ML algorithms, healthcare professionals can analyze patient data to identify areas of improvement in the treatment plan. This can help in providing more personalized and effective care, which can ultimately improve patient outcomes.

Page | 8

It is important to note that the implementation of AI and ML solutions should not be viewed as a replacement for traditional mental healthcare services. Rather, AI and ML tools should be used as a complement to traditional services, to enhance the quality and accessibility of mental healthcare services.

To ensure the successful implementation of AI and ML solutions in mental healthcare services in developing countries, several actions can be taken. Firstly, there is a need to invest in data infrastructure and resources to collect and analyze data. This can be done by governments and international organizations, which can provide funding and technical assistance to developing countries.

Secondly, there is a need to invest in training programs to develop the skills of healthcare professionals in using AI and ML tools. This can be done by universities, training institutes, and professional associations, which can provide training programs and workshops on the use of AI and ML in mental healthcare services. Thirdly, there is a need to raise awareness and acceptance of AI and ML among patients and healthcare professionals. This can be done through education and awareness campaigns, which can highlight the benefits of using AI and ML in mental healthcare services, and address any concerns or misconceptions about the use of these tools. There is a need to ensure that the use of AI and ML in mental healthcare services is done in a responsible and ethical manner. This includes ensuring the privacy and confidentiality of patient data, and avoiding any biases or discrimination in the use of AI and ML algorithms.

The field of natural language processing (NLP) has enormous potential to improve access to mental healthcare services in developing countries. In many of these countries, there is a significant shortage of mental health professionals, and people often face significant barriers when seeking treatment. NLP technologies can help to overcome these barriers by enabling remote consultations, automating certain aspects of diagnosis and treatment, and improving the accuracy and effectiveness of communication between patients and healthcare providers. In this essay, we will explore the various ways in which NLP is being used to improve access to mental healthcare services in developing countries.

One of the key ways in which NLP is being used to improve access to mental healthcare services in developing countries is through the development of chatbots and other conversational agents. These technologies can provide users with a quick and easy way to access mental health information and support, without the need for a human therapist or counsellor. Chatbots can be designed to ask users questions about their symptoms and provide them with advice on how to manage their condition. They can also be programmed to offer cognitive behavioural therapy (CBT) techniques and other evidence-based treatments.

Another way in which NLP is being used to improve access to mental healthcare services in developing countries is through the development of telemedicine platforms. These platforms enable mental health professionals to provide remote consultations to patients who may be unable to attend a physical clinic. This is particularly important in countries where there is a shortage of mental health professionals, or where patients may live in remote or rural areas. Telemedicine platforms can also be used to connect patients with mental health professionals who speak their native language or who have a cultural understanding of their experiences.

NLP can also be used to improve the accuracy and effectiveness of communication between patients and healthcare providers. In many developing countries, there may be language barriers between patients and mental health professionals, which can make it difficult for patients to describe their symptoms and for healthcare providers to understand them. NLP technologies can help to overcome these barriers by providing real-time translation services or by enabling healthcare providers to communicate with patients using simple, non-technical language. This can improve the accuracy of diagnosis and treatment, and ensure that patients receive the care they need.

Furthermore, NLP can be used to automate certain aspects of mental healthcare, such as the identification of high-risk patients and the monitoring of treatment progress. By analysing large datasets of patient records and identifying patterns in language use and behaviour, NLP algorithms can help to identify patients who may be at risk of self-harm or suicide. This can enable mental health professionals to intervene early and provide the necessary support to prevent harm. NLP algorithms can also be used to monitor treatment progress, by analysing patient language and behaviour over time to identify changes in mood or symptoms.

NLP technologies have enormous potential to improve access to mental healthcare services in developing countries. From chatbots and telemedicine platforms to language translation services and automated diagnosis and treatment, there are many ways in which NLP is being used to overcome the barriers that prevent people from accessing the care they need. By leveraging the power of NLP, we can ensure that everyone, regardless of their location or background, has access to the mental healthcare services they need to live healthy and fulfilling lives.

One of the challenges facing the use of NLP in mental healthcare in developing countries is the lack of data. Many NLP algorithms rely on large datasets of patient records and language samples in order to be effective. However, in many developing countries, there is a shortage of mental health professionals and a lack of infrastructure to collect and manage

patient data. This can limit the effectiveness of NLP technologies and make it difficult to develop algorithms that are tailored to the specific needs of local populations.

Another challenge is the potential for bias in NLP algorithms. This is particularly important in mental healthcare, where language use and behaviour can be highly subjective and context-dependent. If NLP algorithms are trained on biased or incomplete datasets, they may produce inaccurate or unfair results. This could have serious consequences for patients, particularly in developing countries where access to mental healthcare is already limited.

To address these challenges, it is important to ensure that NLP algorithms are developed in collaboration with mental health professionals and patients from a diverse range of backgrounds. This can help to ensure that algorithms are tailored to the specific needs of local populations and are free from bias. It is also important to invest in infrastructure and training for mental health professionals in developing countries, in order to improve the quality and quantity of patient data that is available for NLP research. Despite these challenges, the potential benefits of NLP in mental healthcare in developing countries are significant. By leveraging the power of technology, we can help to overcome the barriers that prevent people from accessing the care they need and improve the quality of mental healthcare services around the world.

CONCLUSION

Access to mental healthcare services is a critical issue in developing countries. Mental health conditions are common in these countries, but many individuals do not receive the care they need due to a range of barriers, including stigma, limited resources, and a shortage of mental healthcare providers. Improving access to mental healthcare services is essential to promoting overall health and well-being and reducing the burden of mental illness on individuals, families, and communities in developing countries.

One of the primary barriers to mental healthcare access in developing countries is stigma. Stigma associated with mental health conditions can lead to discrimination and prejudice, and many individuals may feel ashamed or embarrassed to seek help for their mental health challenges. This can lead to delays in seeking treatment or a reluctance to seek help at all. Stigma can also contribute to a lack of investment in mental healthcare services, as mental health conditions may not be seen as a priority by policymakers and funders. To address this issue, it is important to raise awareness about mental health and to reduce the stigma associated with mental illness. This can be achieved through public education campaigns, advocacy, and promoting positive representations of mental health conditions in media and popular culture.

Another barrier to mental healthcare access in developing countries is limited resources. Mental healthcare services can be expensive, and many individuals may not have access to affordable care. In addition, mental healthcare resources may be limited in many areas, particularly in rural or underserved communities. This can make it difficult for individuals to access care, particularly if they do not have the means to travel to another location to

receive treatment. To address this issue, it is important to invest in mental healthcare infrastructure and resources in developing countries. This can include the creation of telehealth programs that allow individuals to receive mental healthcare services remotely and the establishment of mental healthcare clinics in underserved areas.

A shortage of mental healthcare providers is another significant barrier to mental healthcare access in developing countries. There is a significant shortage of mental healthcare providers in many areas, which can make it difficult for individuals to access care. This shortage is particularly acute in developing countries, where there are fewer mental healthcare providers per capita than in developed countries. To address this issue, it is important to invest in mental healthcare provider training programs and to provide incentives for mental healthcare providers to practice in underserved areas.

Cultural competence is another important factor affecting mental healthcare access in developing countries. Mental healthcare providers must be able to provide care that is culturally sensitive and responsive to the needs of diverse populations. This includes individuals from different ethnic and racial backgrounds, individuals with disabilities, and individuals from the LGBTQ+ community. Cultural competence is important because it helps to ensure that individuals receive care that is respectful, responsive, and appropriate to their unique needs and circumstances. To address this issue, it is important to provide cultural competence training to mental healthcare providers and to ensure that mental healthcare services are designed to meet the unique needs of diverse populations.

In addition to these barriers, there are also systemic issues that affect mental healthcare access in developing countries. One of these issues is the lack of integration between mental healthcare services and other healthcare services. Mental healthcare services are often delivered separately from other healthcare services, which can lead to gaps in care and difficulty navigating the healthcare system. To address this issue, it is important to create integrated healthcare systems that provide coordinated and comprehensive care across all aspects of healthcare, including mental healthcare.

Another systemic issue that affects mental healthcare access in developing countries is the lack of investment in mental healthcare research. Research is essential to understanding the causes and effective treatments for mental health conditions, but many developing countries do not have the resources to invest in mental healthcare research. This can limit the development of effective treatments and interventions for mental health conditions in these countries. To address this issue, it is important to invest in mental healthcare research in developing countries and to promote collaborations between mental healthcare researchers and institutions across different countries and regions.

Despite these barriers and systemic issues, there have been some efforts to improve mental healthcare access in developing countries. One approach has been to increase the availability of low-cost or free mental healthcare services. This can be achieved through government-funded programs, non-governmental organizations, and public-private partnerships. For example, in India, the government has established a network of mental healthcare clinics that provide free or low-cost services to individuals with mental health conditions.

Another approach has been to use community-based interventions to promote mental health and prevent mental health conditions. This can include programs that promote mental health literacy, provide social support, and promote healthy lifestyles. For example, in South Africa, a community-based program called "Khuluma" provides mental health education and support to individuals and families affected by HIV/AIDS.

There has also been a growing recognition of the importance of mental healthcare in achieving broader development goals. The United Nations' Sustainable Development Goals include a target to "promote mental health and well-being for all." This recognition has led to increased investment in mental healthcare services and research in many developing countries.

Page | 12

However, there is still much work to be done to improve mental healthcare access in developing countries. One key area of focus is addressing the shortage of mental healthcare providers. This can be achieved through training and education programs that increase the number of mental healthcare providers and promote their practice in underserved areas. In addition, the use of telehealth technologies can help to increase the reach of mental healthcare providers, particularly in remote or underserved areas.

Another important area of focus is addressing the stigma associated with mental health conditions. This can be achieved through public education campaigns, media campaigns, and advocacy efforts. It is also important to promote positive representations of mental health conditions in popular culture and media, to reduce the shame and stigma associated with seeking mental healthcare.

Finally, it is important to recognize the interconnectedness between mental healthcare and broader development goals. Improving mental healthcare access can have significant impacts on economic development, social inclusion, and overall well-being. It is therefore essential to prioritize mental healthcare in development efforts and to invest in mental healthcare infrastructure, research, and services in developing countries.

Access to mental healthcare services is a critical issue in developing countries. Stigma, limited resources, a shortage of mental healthcare providers, cultural competence, and systemic issues all contribute to the challenge of accessing mental healthcare services in these countries. However, there have been efforts to improve mental healthcare access through community-based interventions, low-cost or free mental healthcare services, and increased investment in mental healthcare research and infrastructure. Addressing the barriers to mental healthcare access in developing countries is essential to promoting overall health and well-being and reducing the burden of mental illness on individuals, families, and communities.

REFERENCES

- [1] B. C. Frueh, "Solving Mental Healthcare Access Problems in the Twenty-first Century," *Aust. Psychol.*, vol. 50, no. 4, pp. 304–306, Aug. 2015.

- [2] K. Makiyama *et al.*, “Trends in Healthcare Access in Japan during the First Wave of the COVID-19 Pandemic, up to June 2020,” *Int. J. Environ. Res. Public Health*, vol. 18, no. 6, Mar. 2021.
- [3] D. R. Williams, H. W. Neighbors, and J. S. Jackson, “Racial/ethnic discrimination and health: findings from community studies,” *Am. J. Public Health*, vol. 93, no. 2, pp. 200–208, Feb. 2003.
- [4] B. Giles-Corti, S. Foster, T. Shilton, and R. Falconer, “The co-benefits for health of investing in active transportation,” *N. S. W. Public Health Bull.*, vol. 21, no. 5–6, pp. 122–127, May-Jun 2010.
- [5] R. P. Schumaker, M. A. Veronin, and R. R. Dixit, “Determining Mortality Likelihood of Opioid Drug Combinations using Decision Tree Analysis,” 2022.
- [6] S. Karakolias, C. Kastanioti, M. Theodorou, and N. Polyzos, “Primary care doctors’ assessment of and preferences on their remuneration,” *Inquiry*, vol. 54, p. 46958017692274, Jan. 2017.
- [7] K. Womersley, K. Ripullone, and J. E. Hirst, “Tackling inequality in maternal health: Beyond the postpartum,” *Future Healthc. J.*, vol. 8, no. 1, pp. 31–35, Mar. 2021.
- [8] E. Kim, J. Kim, J. Park, H. Ko, and Y. Kyung, “TinyML-Based Classification in an ECG Monitoring Embedded System,” *CMC-COMPUTERS MATERIALS & CONTINUA*, vol. 75, no. 1, pp. 1751–1764, 2023.
- [9] F. Khayat, L. Teron, and F. Rasoulyan, “COVID-19 and health inequality: the nexus of race, income and mortality in New York City,” *Int. J. Hum. Rights Healthc.*, vol. 15, no. 4, pp. 363–372, Aug. 2022.
- [10] N. Polyzos *et al.*, “Greek National E-Prescribing System: Preliminary Results of a Tool for Rationalizing Pharmaceutical Use and Cost,” *Glob. J. Health Sci.*, vol. 8, no. 10, p. 55711, Oct. 2016.
- [11] K. A. Jablonski and M. F. Guagliardo, “Pediatric appendicitis rupture rate: a national indicator of disparities in healthcare access,” *Popul. Health Metr.*, vol. 3, no. 1, p. 4, May 2005.
- [12] B. Choi, Y. Lee, Y. Kyung, and E. Kim, “ALBERT with Knowledge Graph Encoder Utilizing Semantic Similarity for Commonsense Question Answering,” *arXiv [cs.CL]*, 14-Nov-2022.
- [13] R. Ramalho *et al.*, “Telepsychiatry and healthcare access inequities during the COVID-19 pandemic,” *Asian J. Psychiatr.*, vol. 53, p. 102234, Oct. 2020.
- [14] S. Javanparast, A. Windle, T. Freeman, and F. Baum, “Community Health Worker Programs to Improve Healthcare Access and Equity: Are They Only Relevant to Low- and Middle-Income Countries?,” *Int J Health Policy Manag*, vol. 7, no. 10, pp. 943–954, Oct. 2018.
- [15] M. Mirza *et al.*, “Barriers to healthcare access among refugees with disabilities and chronic health conditions resettled in the US Midwest,” *J. Immigr. Minor. Health*, vol. 16, no. 4, pp. 733–742, Aug. 2014.
- [16] M. S. Winchester and B. King, “Decentralization, healthcare access, and inequality in Mpumalanga, South Africa,” *Health Place*, vol. 51, pp. 200–207, May 2018.
- [17] R. P. Schumaker, M. A. Veronin, T. Rohm, M. Boyett, and R. R. Dixit, “A data driven approach to profile potential SARS-CoV-2 drug interactions using TylerADE,” *J. Int. Technol. Inf. Manag.*, vol. 30, no. 3, pp. 108–142, Jan. 2021.
- [18] P. A. Amoah, J. Edusei, and D. Amuzu, “Social Networks and Health: Understanding the Nuances of Healthcare Access between Urban and Rural Populations,” *Int. J. Environ. Res. Public Health*, vol. 15, no. 5, May 2018.

- [19] M. A. Veronin, R. Dixit, and R. P. Schumaker, "A Decision Tree Analysis of Opioid and Prescription Drug Interactions Leading to Death Using the FAERS Database," in *IIMA/ICITED Joint Conference 2018*, 2018, pp. 67–67.
- [20] Y. Apostolopoulos, S. Sönmez, M. M. Shattell, C. Gonzales, and C. Fehrenbacher, "Health survey of U.S. long-haul truck drivers: work environment, physical health, and healthcare access," *Work*, vol. 46, no. 1, pp. 113–123, Jan. 2013.
- [21] A. Vozikis, A. Panagiotou, and S. Karakolias, "A Tool for Litigation Risk Analysis for Medical Liability Cases," *HAPScPBS*, vol. 2, no. 2, pp. 268–277, Dec. 2021.
- [22] S. Lim, P. Y. Chan, S. Walters, G. Culp, M. Huynh, and L. H. Gould, "Impact of residential displacement on healthcare access and mental health among original residents of gentrifying neighborhoods in New York City," *PLoS One*, vol. 12, no. 12, p. e0190139, Dec. 2017.
- [23] J. Wagner, G. Burke, T. Kuoch, M. Scully, S. Armeli, and T. V. Rajan, "Trauma, healthcare access, and health outcomes among Southeast Asian refugees in Connecticut," *J. Immigr. Minor. Health*, vol. 15, no. 6, pp. 1065–1072, Dec. 2013.
- [24] N. Polyzos, S. Karakolias, G. Mavridoglou, P. Gkorezis, and C. Zilidis, "Current and future insight into human resources for health in Greece," *Open J. Soc. Sci.*, vol. 03, no. 05, pp. 5–14, 2015.
- [25] M. E. Cupples, P. M. Hart, A. Johnston, and A. J. Jackson, "Improving healthcare access for people with visual impairment and blindness," *BMJ*, vol. 344, p. e542, Jan. 2012.
- [26] E. Kim, Y. Lee, J. Choi, B. Yoo, K. J. Chae, and C. H. Lee, "Machine Learning-based Prediction of Relative Regional Air Volume Change from Healthy Human Lung CTs," *KSII Transactions on Internet & Information Systems*, vol. 17, no. 2, 2023.
- [27] S. Karakolias and C. Kastanioti, "Application of an organizational assessment tool of primary health care," *Arch Hell Med*, vol. 35, pp. 497–505, 2018.
- [28] J. Gelatt, "Immigration status and the healthcare access and health of children of immigrants," *Soc. Sci. Q.*, vol. 97, no. 3, pp. 540–554, Sep. 2016.
- [29] S. Karakolias and N. Polyzos, "Application and assessment of a financial distress projection model in private general clinics," *Archives of Hellenic Medicine/Arheia Ellenikes Iatrikes*, vol. 32, no. 4, 2015.
- [30] J.-H. Tang, Y.-H. Chiu, P.-H. Chiang, M.-D. Su, and T.-C. Chan, "A flow-based statistical model integrating spatial and nonspatial dimensions to measure healthcare access," *Health Place*, vol. 47, pp. 126–138, Sep. 2017.
- [31] M. B. Adugna, F. Nabbouh, S. Shehata, and S. Ghahari, "Barriers and facilitators to healthcare access for children with disabilities in low and middle income sub-Saharan African countries: a scoping review," *BMC Health Serv. Res.*, vol. 20, no. 1, p. 15, Jan. 2020.
- [32] H. Elath, R. R. Dixit, R. P. Schumaker, and M. A. Veronin, "Predicting Deadly Drug Combinations through a Machine Learning Approach," in *PACIS*, 2018, p. 177.
- [33] S. E. Karakolias and N. M. Polyzos, "The newly established unified healthcare fund (EOPYY): current situation and proposed structural changes, towards an upgraded model of primary health care, in Greece," *Health*, vol. 2014, 2014.
- [34] M. Gunst *et al.*, "Healthcare access for refugees in Greece: Challenges and opportunities," *Health Policy*, vol. 123, no. 9, pp. 818–824, Sep. 2019.
- [35] E. Kim and Y. Kyung, "Factors Affecting the Adoption Intention of New Electronic Authentication Services: A Convergent Model Approach of VAM, PMT, and TPB," *IEEE Access*, vol. 11, pp. 13859–13876, 2023.
- [36] K. Macapagal, R. Bhatia, and G. J. Greene, "Differences in Healthcare Access, Use, and Experiences Within a Community Sample of Racially Diverse Lesbian, Gay,

- Bisexual, Transgender, and Questioning Emerging Adults,” *LGBT Health*, vol. 3, no. 6, pp. 434–442, Dec. 2016.
- [37] N. Polyzos *et al.*, “The introduction of Greek Central Health Fund: Has the reform met its goal in the sector of Primary Health Care or is there a new model needed?,” *BMC Health Serv. Res.*, vol. 14, p. 583, Nov. 2014.
- [38] M. P. Rivera *et al.*, “Addressing Disparities in Lung Cancer Screening Eligibility and Healthcare Access. An Official American Thoracic Society Statement,” *Am. J. Respir. Crit. Care Med.*, vol. 202, no. 7, pp. e95–e112, Oct. 2020.
- [39] M. J. Haenssgen and P. Ariana, “Healthcare access: A sequence-sensitive approach,” *SSM Popul Health*, vol. 3, pp. 37–47, Dec. 2017.
- [40] N. Fullman *et al.*, “Measuring performance on the Healthcare Access and Quality Index for 195 countries and territories and selected subnational locations: a systematic analysis from the Global Burden of Disease Study 2016,” *Lancet*, vol. 391, no. 10136, pp. 2236–2271, 2018.
- [41] A. S. W. Chan and Others, “Safe is not enough: better schools for LGBTQ students (youth development and education series)[Book review],” 2021.
- [42] D. Meshvara, “Mental health and mental health care in Asia,” *World Psychiatry*, vol. 1, no. 2, pp. 118–120, Jun. 2002.
- [43] A. S. W. Chan, J. M. C. Ho, and P. M. K. Tang, “Cancer and the LGBT Community: Cancer and the LGBT Community (2015th ed.), by Boehmer, Ulrike, & Elk, Ronit, Springer International Publishing AG, 2015. https ...,” 2021.
- [44] A. S. W. Chan, J. S. F. Li, J. M. C. Ho, H. L. Tam, and W. L. Hsu, “The systematic review and meta-analysis of Chronic Inflammation and Fibrosis in HIV/AIDS and Cancer: Impacts of Psychological Wellbeing among ...,” *Frontiers in Public Health*.
- [45] A. S. W. Chan *et al.*, “Impacts of psychological wellbeing with HIV/AIDS and cancer among sexual and gender minorities: A systematic review and meta-analysis,” *Front Public Health*, vol. 10, p. 912980, Nov. 2022.
- [46] A. S. W. Chan, P. M. K. Tang, and E. Yan, “Chemsex and its risk factors associated with human immunodeficiency virus among men who have sex with men in Hong Kong,” *World Journal of Virology*, 2022.
- [47] A. S. W. Chan, J. M. C. Ho, H. L. Tam, W. L. Hsu, and P. M. K. Tang, “COVID-19, SARS, and MERS: the risk factor associated with depression and its impact on psychological well-being among sexual moralities,” 2022.
- [48] A. S. W. Chan, “Book Review: Safe Is Not Enough: Better Schools for LGBTQ Students (Youth Development and Education Series),” 2021.
- [49] A. S. W. Chan and P. M. K. Tang, “Application of Novel Psychoactive Substances: Chemsex and HIV/AIDS Policies Among Men Who Have Sex With Men in Hong Kong,” *Front. Psychiatry*, vol. 12, p. 680252, Jul. 2021.
- [50] E. Kim *et al.*, “SHOMY: Detection of Small Hazardous Objects using the You Only Look Once Algorithm,” *KSII Transactions on Internet & Information Systems*, vol. 16, no. 8, 2022.
- [51] R. Kakuma *et al.*, “Human resources for mental health care: current situation and strategies for action,” *Lancet*, vol. 378, no. 9803, pp. 1654–1663, Nov. 2011.
- [52] A. S. W. Chan, “Book review: the Educator’s guide to LGBT+ inclusion: a practical resource for K-12 teachers, administrators, and school support staff,” 2021.
- [53] A. S. W. Chan, “Book review: the deviant’s war: the homosexual vs. the United States of America,” 2021.
- [54] A. S. W. Chan, J. M. C. Ho, H. L. Tam, and P. M. K. Tang, “Book review: successful aging: a neuroscientist explores the power and potential of our lives,” *Front. Psychol.*, 2021.

- [55] V. Patel, *Where There Is No Psychiatrist: A mental health care manual*. London, England: Gaskell, 2003.
- [56] A. S. W. Chan, D. Wu, I. P. Y. Lo, J. M. C. Ho, and E. Yan, “Diversity and Inclusion: Impacts on Psychological Wellbeing Among Lesbian, Gay, Bisexual, Transgender, and Queer Communities,” *Front. Psychol.*, vol. 13, p. 726343, Apr. 2022.
- [57] A. S. W. Chan, I. P. Y. Lo, and E. Yan, “Health and Social Inclusion: The Impact of Psychological Well-Being and Suicide Attempts Among Older Men Who Have Sex With Men,” *Am. J. Mens. Health*, vol. 16, no. 5, p. 15579883221120984, Sep-Oct 2022.
- [58] E. Kim, M. Kim, and Y. Kyung, “A Case Study of Digital Transformation: Focusing on the Financial Sector in South Korea and Overseas,” *Asia Pacific Journal of Information Systems*, vol. 32, no. 3, pp. 537–563, 2022.
- [59] A. S. W. Chan, “Book review: the gay revolution: the story of the struggle,” 2021.
- [60] A. S. W. Chan, J. M. C. Ho, J. S. F. Li, and H. L. Tam, “Impacts of COVID-19 pandemic on psychological well-being of older chronic kidney disease patients,” *Frontiers in Medicine*, 2021.
- [61] H. Remschmidt and M. Belfer, “Mental health care for children and adolescents worldwide: a review,” *World Psychiatry*, vol. 4, no. 3, pp. 147–153, Oct. 2005.
- [62] V. W. Xue *et al.*, “USMB-shMincle: a virus-free gene therapy for blocking M1/M2 polarization of tumor-associated macrophages,” *Mol Ther Oncolytics*, vol. 23, pp. 26–37, Dec. 2021.
- [63] E.-C. Kim, E.-Y. Kim, H.-C. Lee, and B.-J. Yoo, “The Details and Outlook of Three Data Acts Amendment in South Korea: With a Focus on the Changes of Domestic Financial and Data Industry,” *Informatization Policy*, vol. 28, no. 3, pp. 49–72, 2021.
- [64] J. Y. F. Chung, M. K. K. Chan, and P. C. T. Tang, “AANG: A natural compound formula for overcoming multidrug resistance via synergistic rebalancing the TGF- β /Smad signalling in hepatocellular carcinoma,” *Journal of Cellular*, 2021.
- [65] H. L. Tam, L. Y. L. Leung, and E. M. L. Wong, “Integration of text messaging interventions into hypertension management among older adults: a systematic review and meta-analysis,” *on Evidence-Based ...*, 2022.
- [66] G. Rojas, V. Martínez, P. Martínez, P. Franco, and Á. Jiménez-Molina, “Improving Mental Health Care in Developing Countries Through Digital Technologies: A Mini Narrative Review of the Chilean Case,” *Front Public Health*, vol. 7, p. 391, Dec. 2019.
- [67] R. Giel *et al.*, “The burden of mental illness on the family,” *Acta Psychiatr. Scand.*, vol. 68, no. 3, pp. 186–201, Sep. 1983.
- [68] A. Javed, M. N. S. Khan, A. Nasar, and A. Rasheed, “Mental healthcare in Pakistan,” *Taiwanese Journal of Psychiatry*, vol. 34, no. 1, p. 6, Jan. 2020.
- [69] M. R. Gutiérrez-Colosía *et al.*, “Standard comparison of local mental health care systems in eight European countries,” *Epidemiol. Psychiatr. Sci.*, vol. 28, no. 2, pp. 210–223, Apr. 2019.
- [70] R. Kohn, S. Saxena, I. Levav, and B. Saraceno, “The treatment gap in mental health care,” *Bull. World Health Organ.*, vol. 82, no. 11, pp. 858–866, Nov. 2004.
- [71] World Health Organization(WHO), *Mental health care in developing countries: A critical appraisal of research findings*. Genève, Switzerland: World Health Organization, 1984.